


 CrossMark
 click for updates

 Cite this: *RSC Adv.*, 2015, 5, 84657

 Received 17th August 2015
 Accepted 30th September 2015

DOI: 10.1039/c5ra16548b

www.rsc.org/advances

Silver-catalyzed direct spirocyclization of alkynes with thiophenols: a simple and facile approach to 3-thioazaspiro[4,5]trienones†


Huanhuan Cui, Wei Wei,* Daoshan Yang, Jimei Zhang, Zhihong Xu, Jiangwei Wen and Hua Wang*

A new and convenient silver-catalyzed direct oxidative spirocyclization of alkynes with thiophenols is described. This methodology provides a simple and practical approach to various 3-thioazaspiro[4,5]-trienones in moderate to good yields with high atom efficiency and excellent functional group tolerance.

As a highly important class of compounds, sulfur-containing molecules exhibit a wide range of applications in organic synthesis,^{1,2} medicinal chemistry,³ and materials science.⁴ Consequently, numerous efforts have been made to explore new efficient methods for introduction of the sulfur group into organic molecules in the synthetic community. Among many of the synthetic transformations, the difunctionalization of C–C unsaturated bonds with thiolation agents has recently attracted great interests of chemists due to it could provide rapid and concise access to various sulfur-containing compounds.^{5–10} Over the past decades, many transition-metal-catalyzed or metal-free difunctionalizations of alkenes for constructing sulfur-containing compounds have been developed,^{5–8} such as alkoxythiolation,⁵ hydroxythiolation,⁶ acetoxythiolation,⁷ and sulfamination.⁸ Nevertheless, to date, only a few strategies for the fabrication of sulfur compounds has been exploited through the difunctionalization of alkynes.^{9,10} Recently, Li *et al.* reported Mn(OAc)₂-catalyzed difunctionalization of alkynes with thiophenols leading to benzothiophenes.^{10a} Zou and co-workers reported oxidative radical oxythiolation of alkynes with thiophenols to access α -thioaldehydes.^{10b} Yi also described iodothiolation of alkynes with sodium arenesulfonates for the construction of β -iodoalkenyl sulfides under metal-free conditions.^{10c} Although some remarkable progress has been made in this field, the development of new, convenient, efficient and atom-economical protocols through difunctionalization of alkynes to assembly structurally diverse and complex sulfur-containing chemical frameworks still remains a highly desirable but challenging task in the modern organic chemistry.

As part of our continued interest in the difunctionalization of alkynes,¹¹ we herein report a new silver-catalyzed oxidative spirocyclization of *N*-arylpropiolamides with thiophenols leading to 3-thioazaspiro[4,5]trienones (eqn (1)). Azaspiro[4,5]-trienone is the common core structure in many natural products and pharmaceuticals as well as a versatile building block in organic synthesis.^{12,13} Recently, several strategies based on the difunctionalization of alkynes has been applied successfully for the construction of various functionalized azaspiro[4,5]-trienones.^{11a,14} Through this methodology, some additional functional groups could be introduced into the azaspiro[4,5]-trienone framework. Very recently, a report on the use of the reagent combination of *N*-(*p*-methoxyaryl)propiolamides, disulfides, CuCl₂, O₂, and H₂O in DMF at 100 °C for the synthesis of 3-thioazaspiro[4,5]trienones was disclosed by Li and Song.¹⁵ However, this method is restricted with the use of the preformed disulfides and arylalkynes bearing the *para*-methoxy substituent on the aryl ring. Moreover, the mechanism for the *ipso*-cyclization method was not studied in detail.

The present protocol offer a convenient and efficient route to a series of biologically important 3-thioazaspiro[4,5]trienones *via* direct silver-catalyzed *ipso*-cyclization of *para*-unsubstituted arylalkynes with commercial available thiophenols, in which the C–S, C–C, and C=O bonds can be sequential formed in a single operation (eqn (1)). Preliminary mechanistic studies suggested that the carbonyl oxygen atom of 3-thioazaspiro[4,5]-trienones originated from the water and this reaction might involve a radical process.




The Key Laboratory of Life-Organic Analysis and Key Laboratory of Pharmaceutical Intermediates and Analysis of Natural Medicine, School of Chemistry and Chemical Engineering, Qufu Normal University, Qufu 273165, Shandong, China. E-mail: weiweiqfnu@163.com; huawang_qfnu@126.com; Web: <http://wang.qfnu.edu.cn>

† Electronic supplementary information (ESI) available: Experimental details. See DOI: 10.1039/c5ra16548b

In our initial studies, the reaction of *N*-methyl-*N*,3-diphenylpropiolamide **1a** with 4-methylbenzenethiol **2a** was performed to examine the catalytic activity of various catalysts including Cu, Pd, Ni, Fe, Ag, and iodized salts in the presence of H₂O (3 equiv.) in CH₃CN under air. As shown in Table 1, among those catalysts tested (Table 1, entries 1–11), silver salts especially AgCl was found to be the best catalyst to catalyze the formation of product **3a** (Table 1, entry 8). Among the solvents examined, 1,4-dioxane was demonstrated to be more effective than the others such as CH₃CN, DME, DCE, THF, toluene, DMF, and DMSO (Table 1, entries 8, 12–18). In addition, the reaction efficiency was obviously low with the decreasing of AgCl loading (Table 1, entries 19 and 20). We found that the reaction temperature played an important role in this reaction (Table 1, entries 12, 21–23). The desired product was obtained in only 14% yield when the model reaction was carried out at room temperature (Table 1, entry 21), and the reaction at 80 °C gave the best results. After a series of detailed investigations, the best yield of **3a** (76%) was obtained by employing **1a** (0.25 mmol), **2a** (0.5 mmol), AgCl (10 mol%), and H₂O (3 equiv.) in 1,4-dioxane at 80 °C (Table 1, entry 12).

Upon optimization of the reaction conditions, the scope of this new spirocyclization reaction was evaluated, with some


Table 1 Optimization of the reaction conditions^a



| Entry | Catalyst | Solvent | Yield ^b (%) |
|-------|--------------------------------------|--------------------|------------------------|
| 1 | CuI | CH ₃ CN | <10 |
| 2 | Pd(OAc) ₂ | CH ₃ CN | 30 |
| 3 | NiCl ₂ | CH ₃ CN | 33 |
| 4 | FeCl ₃ ·6H ₂ O | CH ₃ CN | <10 |
| 5 | Ag ₂ O | CH ₃ CN | 24 |
| 6 | AgOAc | CH ₃ CN | 42 |
| 7 | Ag ₂ CO ₃ | CH ₃ CN | <10 |
| 8 | AgCl | CH ₃ CN | 67 |
| 9 | AgNO ₃ | CH ₃ CN | 65 |
| 10 | KI | CH ₃ CN | 23 |
| 11 | TBAI | CH ₃ CN | 28 |
| 12 | AgCl | 1,4-Dioxane | 76 |
| 13 | AgCl | DME | 65 |
| 14 | AgCl | DCE | 64 |
| 15 | AgCl | THF | 40 |
| 16 | AgCl | Toluene | 68 |
| 17 | AgCl | DMF | 42 |
| 18 | AgCl | DMSO | 36 |
| 19 | AgCl | 1,4-Dioxane | 58 ^c |
| 20 | AgCl | 1,4-Dioxane | 46 ^d |
| 21 | AgCl | 1,4-Dioxane | 14 ^e |
| 22 | AgCl | 1,4-Dioxane | 66 ^f |
| 23 | AgCl | 1,4-Dioxane | 69 ^g |

^a Reaction conditions: **1a** (0.25 mmol), **2a** (0.5 mmol), catalyst (10 mol%), H₂O (3 equiv.), anhydrous solvent (2 mL), 80 °C, 14 h, air. DCE: 1,2-dichloroethane; DME: 1,2-dimethoxyethane. ^b Isolated yields based on **1a**. ^c AgCl (5 mol%). ^d AgCl (2 mol%). ^e 25 °C. ^f 60 °C. ^g 100 °C.


Table 2 Results for the reaction of the spirocyclization of alkynes with thiophenols^{ab}







^a Reaction conditions: **1** (0.25 mmol), **2** (0.5 mmol), AgCl (10 mol%), H₂O (3 equiv.), 1,4-dioxane (2 mL), 10–24 h, 80 °C, under air. ^b Isolated yields based on **1**.

results summarized in Table 2. Firstly, the effects of the substituent on the alkynyl moiety were investigated. Arylalkynes bearing both of the electron-donating and electron-withdrawing groups on the aromatic moieties were tolerated in this reaction to give the corresponding products in good yields (**3a–3e**). As expected, alkylalkyne was also suitable for this reaction, but leading to the desired product in the relatively lower yield (**3f**). Subsequently, *N*-arylpropiolamides with various substitution patterns at the aniline moieties were examined. The *ortho*- or

meta-position of the aniline moieties were compatible with this reaction, with the desired products obtained in moderate to good yields (**3g–3n**). Notably, various halogen groups were consistent with the optimized conditions, thereby facilitating further transformations (**3i–3l**). Interestingly, the *para*-position substituted *N*-arylpropiolamides could give the corresponding product **3a** in moderate to good yields by releasing the *para*-substituents (*p*-MeO-, *p*-F, and *p*-Br) (eqn (2)). Meantime, naphthyl moiety could also be used in the reactions to give the expected product **3o** in 60% yield. In addition, the reaction could also proceed well by using various thiophenols with an electron-donating group (Me or MeO) or an electron-withdrawing group (Cl or Br) on the aromatic ring to give the corresponding products in moderate to good yields (**3p–3v**). Nevertheless, none of the desired product was obtained when changing the *N*-Me group to the *N*-H or *N*-Ph group (**3w** and **3x**), which might be caused by the electronic effect.




Several control experiments were conducted to obtain some insights into this reaction (eqn (3)–(6)). Azaspiro[4,5]trienone **1a'** was not detected when *N*-methyl-*N*,3-diphenylpropiolamide **1a** was performed dependently under the standard conditions, indicating that azaspiro[4,5]trienone **1a'** might not be the key intermediate in this reaction (eqn (3)). Furthermore, this reaction was completely inhibited when TEMPO (2,2,6,6-tetramethyl-1-piperidinyloxy, a well known radical-capturing species) was added into reaction system under the standard condition and TEMPO-trapped complex (*p*-MePhS-TEMPO) was detected by Lc-Ms analysis (see ESI[†]), which suggested that ArS radical might involve in the present reaction system and this transformation should proceed through a radical pathway (eqn (4)). Moreover, ¹⁸O atom-labelling experiment demonstrated that ¹⁸O atom could be incorporated into the corresponding product **3a** when the reaction of **1a** with **2a** was carried out in the presence of H₂¹⁸O (eqn (5)). This result showed that the oxygen atom of carbonyl group originated from water. In addition, the intermolecular kinetic isotope effect (KIE) experiment were carried out with the deuterium labeled substrates (eqn (6)). No kinetic isotope effects ($k_H/k_D = 1.0$) was observed, which indicated C–H bond cleavage might not be the rate-determining step that was similar to mechanisms proposed in previous reports.^{11a,14}



Based on the above experimental results and previous reports,^{11a,14–18} a postulated reaction pathway was proposed as shown in Scheme 1. Initially, the thiyl radical **4** was generated from thiophenol **2** in the presence of silver salt under air.¹⁶ Then, the addition of thiyl radical **4** to the triple bond of arylpropiolamide **1** gave vinyl radical **5**. Subsequently, the intramolecular *spiro*-cyclization of vinyl radical **5** with an aryl ring would lead to the formation of the radical intermediate **6**. Next, the corresponding cyclohexadienyl cation **7** was produced from intermediate **6** *via* a single-electron-transfer process. Finally, the nucleophilic addition of H₂O to cation **7** afforded intermediate **8**,^{11a,14i,17} which was further oxidized to form the desired product **3**.^{11a,14i,18}

In conclusion, a simple and efficient protocol has been successfully developed for the construction of 3-thioazaspiro[4,5]trienones *via* silver-catalyzed oxidative spirocyclization of *N*-arylpropiolamides with thiophenols. Preliminary mechanistic studies indicated that this reaction might involve a radical process and the carbonyl oxygen atom of 3-thioazaspiro[4,5]trienones originated from the water. This simple reaction system is expected to expand the potential applications



Scheme 1 Postulated reaction pathway.

of functionalized azaspiro[4,5]trienones in the synthetic and pharmaceutical chemistry.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (No. 21302109, 21302110, and 21375075), the Taishan Scholar Foundation of Shandong Province, the Natural Science Foundation of Shandong Province (ZR2015JL004).

Notes and references

- 1 Selective examples see: (a) I. P. Beletskaya and V. P. Ananikov, *Eur. J. Org. Chem.*, 2007, 3431; (b) M. D. McReynolds, J. M. Dougherty and P. R. Hanson, *Chem. Rev.*, 2004, **104**, 2239; (c) N. V. Zyk, E. K. Beloglazkina, M. A. Belova and N. S. Dubinina, *Russ. Chem. Rev.*, 2003, **72**, 769; (d) A. Y. Sizov, A. N. Kovregin and A. F. Ermolov, *Russ. Chem. Rev.*, 2003, **72**, 357; (e) T. do and T. Mitsudo, *Chem. Rev.*, 2000, **100**, 3205; (f) A. Mangini, *Sulfur Rep.*, 1987, **7**, 313; (g) E. Block, *Reactions of Organosulfur Compounds*, Academic Press, New York, 1978.
- 2 (a) R. Sarma, N. Rajesh and D. Prajapati, *Chem. Commun.*, 2012, **48**, 4014; (b) N. Muraoka, M. Mineno, K. Itami and J.-I. Yoshida, *J. Org. Chem.*, 2005, **70**, 6933; (c) W. H. Pearson, I. Y. Lee, Y. Mi and P. Stoy, *J. Org. Chem.*, 2004, **69**, 9109; (d) S. Farhat and I. Marek, *Angew. Chem., Int. Ed.*, 2002, **41**, 1410; (e) M. Su, W. Yu and Z. Jin, *Tetrahedron Lett.*, 2001, **42**, 3771; (f) H. Mizuno, K. Domon, K. Asuya, K. Tanino and I. Kuwajima, *J. Org. Chem.*, 1999, **64**, 2648; (g) D. F. Andres, U. Dietrich, E. G. Laurent and B. S. Marquet, *Tetrahedron*, 1997, **53**, 647; (h) J.-E. Baekvall and A. Ericsson, *J. Org. Chem.*, 1994, **59**, 5850.
- 3 (a) J. M. Paris, J. C. Barriere, C. Smith and P. E. Bost, in *Recent Progress in the Chemical Synthesis of Antibiotics*, ed. G. Lucas and M. Ohno, Springer-Verlag, Berlin, 1990, p. 183; (b) M. Di Giambattista, G. Chinali and C. J. Cocito, *J. Antimicrob. Chemother.*, 1989, **24**, 485; (c) C. Cocito, *Microbiol. Rev.*, 1979, **43**, 145.
- 4 (a) C. Lauterbach and J. Fabian, *Eur. J. Inorg. Chem.*, 1999, 1995; (b) E. G. Hope and W. Levason, *Coord. Chem. Rev.*, 1993, **122**, 109; (c) P. I. Clemenson, *Coord. Chem. Rev.*, 1990, **106**, 171; (d) R. P. Baumann, K. Ishiguro, P. G. Penketh, K. Shyam, R. Zhu and A. C. Sartorelli, *Biochem. Pharmacol.*, 2011, **81**, 1201.
- 5 (a) H. Guan, H. Wang, D. Huang and Y. Shi, *Tetrahedron*, 2012, **68**, 2728; (b) H. Wang, D. Huang, D. Cheng, L. Li and Y. Shi, *Org. Lett.*, 2011, **13**, 1650; (c) X. Gao, X. Pan, J. Gao, H. Jiang, G. Yuan and Y. Li, *Org. Lett.*, 2015, **17**, 1038; (d) F.-L. Yang, F.-X. Wang, T.-T. Wang, Y.-J. Wang and S.-K. Tian, *Chem. Commun.*, 2014, **50**, 2111; (e) J. Yu, C. Gao, Z. Song, H. Yang and H. Fu, *Org. Biomol. Chem.*, 2015, **13**, 4846; (f) A. Vieira, J. B. Azeredo, M. Godoi, C. Santi, E. N. da Silva Júnior and A. L. Braga, *J. Org. Chem.*, 2015, **80**, 2120.
- 6 (a) A. Bewick, J. M. Mellor, D. Milano and W. M. Owton, *J. Chem. Soc., Perkin Trans. 1*, 1985, 1045; (b) L. D. S. Yadav and C. Awasthi, *Tetrahedron Lett.*, 2009, **50**, 3801; (c) B. Movassagh and M. Navidi, *Tetrahedron Lett.*, 2008, **49**, 6712; (d) K. Surendra, N. S. Krishnaveni, R. Sridhar and K. R. Rao, *J. Org. Chem.*, 2006, **71**, 5819; (e) D. Ahmed Kamal and R. Rajasekhar Reddy, *J. Mol. Catal. A: Chem.*, 2007, **272**, 26; (f) T. Keshari, V. K. Yadav, V. P. Srivastava and L. D. S. Yadav, *Green Chem.*, 2014, **16**, 3986; (g) H. Xi, B. Deng, Z. Zong, S. Lu and Z. Li, *Org. Lett.*, 2015, **17**, 1180; (h) V. K. Yadav, V. P. Srivastava and L. D. S. Yadav, *Tetrahedron Lett.*, 2015, **56**, 2892.
- 7 (a) N. Taniguchi, *J. Org. Chem.*, 2006, **71**, 7874; (b) C. Muangkaew, P. Katrun, P. Kanchanarugee, M. Pohmakotr, V. Reutrakul, D. Soorukram, T. Jaipetch and C. Kuhakarn, *Tetrahedron*, 2013, **69**, 8847.
- 8 (a) L. Li, H. Wang, D. Huang and Y. Shi, *Tetrahedron*, 2012, **68**, 9853; (b) L. Li, Z. Li, D. Huang, H. Wang and Y. Shi, *RSC Adv.*, 2013, **3**, 4523–4525.
- 9 (a) N. Taniguchi, *Tetrahedron*, 2009, **65**, 2782; (b) N. Taniguchi, *Synlett*, 2008, 849; (c) X. Zeng, L. Ilies and E. Nakamura, *Org. Lett.*, 2012, **14**, 954; (d) M. Iwasaki, T. Fujii, A. Yamamoto, K. Nakajima and Y. Nishihara, *Chem.-Asian J.*, 2014, **9**, 58; (e) M. Iwasaki, T. Fujii, K. Nakajima and Y. Nishihara, *Angew. Chem., Int. Ed.*, 2014, **53**, 13880; (f) L. Zhong, P. R. Savoie, A. S. Filatov and J. T. Welch, *Angew. Chem., Int. Ed.*, 2014, **53**, 526; (g) X. Zhao, L. Zhang, X. Lu, T. Li and K. Lu, *J. Org. Chem.*, 2015, **80**, 2918.
- 10 (a) K. Liu, F. Jia, H. Xi, Y. Li, X. Zheng, Q. Guo, B. Shen and Z. Li, *Org. Lett.*, 2013, **15**, 2026; (b) S.-F. Zhou, X.-Q. Pan, Z.-H. Zhou, A. Shoberu, P.-Z. Zhang and J.-P. Zou, *J. Org. Chem.*, 2015, **80**, 5348; (c) Y. Lin, G. Lu, C. Cai and W. Yi, *Org. Lett.*, 2015, **17**, 3310.
- 11 (a) J. Wen, W. Wei, S. Xue, D. Yang, Y. Lou, C. Gao and H. Wang, *J. Org. Chem.*, 2015, **80**, 4966; (b) W. Wei, J. Wen, D. Yang, H. Jing, J. You and H. Wang, *RSC Adv.*, 2015, **5**, 4416; (c) D. Yang, K. Yan, W. Wei, L. Tian, Q. Li, J. You and H. Wang, *RSC Adv.*, 2014, **4**, 48547; (d) W. Wei, J. Wen, D. Yang, M. Guo, Y. Wang, J. You and H. Wang, *Chem. Commun.*, 2015, **51**, 768.
- 12 (a) M. Sannigrahi, *Tetrahedron*, 1999, **55**, 9007; (b) Y. Yang, F. Chang and Y. Wu, *Helv. Chim. Acta*, 2004, **87**, 1392; (c) S. Kotha, A. C. Deb, K. Lahiri and E. Manivannan, *Synthesis*, 2009, 165; (d) S. Rosenberg and R. Leino, *Synthesis*, 2009, 2651; (e) E. M. Antunes, B. R. Copp, M. T. Davies-Coleman and T. Samaai, *Nat. Prod. Rep.*, 2005, **22**, 62; (f) E. Gravel and E. Poupon, *Nat. Prod. Rep.*, 2010, **27**, 32; (g) Y.-S. Cai, Y.-W. Guo and K. Krohn, *Nat. Prod. Rep.*, 2010, **27**, 1840.
- 13 (a) I. Hermecz, A. Santa-Csutor, C. Gönczi, G. Héja, E. Csikós, K. Simon, A. Smelkó-Esek and B. Podányi, *Pure Appl. Chem.*, 2001, **71**, 1401; (b) J. Halász, B. Podányi, A. Santa-Csutor, Z. Böcskei, K. Simon, M. Hanusz and I. Hermecz, *J. Mol. Struct.*, 2003, 187; (c) S. T. Roche and J. A. Porco Jr, *Angew. Chem., Int. Ed.*, 2011, **50**, 4068; (d) C.-X. Zhuo, W. Zhang and S.-L. You, *Angew. Chem., Int. Ed.*, 2012, **51**, 12662.

- 14 Representative examples see: (a) X. Zhang and R. C. Larock, *J. Am. Chem. Soc.*, 2005, **127**, 12230; (b) B.-X. Tang, D.-J. Tang, S. Tang, Q.-F. Yu, Y.-H. Zhang, Y. Liang, P. Zhong and J.-H. Li, *Org. Lett.*, 2008, **10**, 1063; (c) B. Godoi, R. F. Schumacher and G. Zeni, *Chem. Rev.*, 2011, **111**, 2937; (d) B.-X. Tang, Y.-H. Zhang, R.-J. Song, D.-J. Tang, G.-B. Deng, Z.-Q. Wang, Y.-X. Xie, Y.-Z. Xia and J.-H. Li, *J. Org. Chem.*, 2012, **77**, 2837; (e) B.-X. Tang, Q. Yin, R.-Y. Tang and J.-H. Li, *J. Org. Chem.*, 2008, **73**, 9008; (f) W.-T. Wei, R.-J. Song, X.-H. Ouyang, Y. Li, H.-B. Li and J.-H. Li, *Org. Chem. Front.*, 2014, **1**, 484; (g) X.-H. Ouyang, R.-J. Song, Y. Li, B. Liu and J.-H. Li, *J. Org. Chem.*, 2014, **79**, 4582; (h) L.-J. Wang, A.-Q. Wang, Y. Xia, X.-X. Wu, X.-Y. Liu and Y.-M. Liang, *Chem. Commun.*, 2014, **50**, 13998; (i) X.-H. Yang, X.-H. Ouyang, W.-T. Wei, R.-J. Song and J.-H. Li, *Adv. Synth. Catal.*, 2015, **357**, 1161.
- 15 P.-C. Qian, Y. Liu, R.-J. Song, J.-N. Xiang and J.-H. Li, *Synlett*, 2015, **26**, 1213.
- 16 A. K. Singh, R. Chawla, T. Keshari, V. K. Yadav and L. D. S. Yadav, *Org. Biomol. Chem.*, 2014, **12**, 8550.
- 17 (a) H. J. Kim, J. Kim, S. H. Cho and S. Chang, *J. Am. Chem. Soc.*, 2011, **133**, 16382; (b) Y. Yan, Y. Zhang, Z. Zha and Z. Y. Wang, *Org. Lett.*, 2013, **15**, 2274.
- 18 (a) B.-X. Tang, Y.-H. Zhang, R.-J. Song, D.-J. Tang, G.-B. Deng, Z.-Q. Wang, Y.-X. Xie, Y.-Z. Xia and J.-H. Li, *J. Org. Chem.*, 2012, **77**, 2837; (b) G. Han, Q. Wang, Y. Liu and Q. Wang, *Org. Lett.*, 2014, **16**, 5914.