

### Copper-catalyzed highly selective direct hydrosulfonylation of alkynes with arylsulfonic acids leading to vinyl sulfones

Cite this: *Org. Biomol. Chem.*, 2014, **12**, 1861

Received 17th December 2013,

Accepted 21st January 2014


DOI: 10.1039/c3ob42522c

www.rsc.org/obc

Wei Wei,<sup>a,b</sup> Jinli Li,<sup>a,b</sup> Daoshan Yang,<sup>a,b</sup> Jiangwei Wen,<sup>a,b</sup> Yueting Jiao,<sup>a,b</sup>  
Jinmao You<sup>a,b,c</sup> and Hua Wang<sup>\*a,b</sup>

**A novel Cu-catalyzed direct hydrosulfonylation of alkynes with arylsulfonic acids for the synthesis of (*E*)-vinyl sulfones has been realized under mild conditions with 100% atom efficiency. The present protocol provides an attractive approach to various vinyl sulfones in good to excellent yields, with the advantages of operation simplicity, atom economy, and high stereo- and regioselectivities.**

Transition-metal-catalyzed direct addition of H-heteroatom compounds to carbon-carbon unsaturated triple bonds is one of the most straightforward and powerful tools for the construction of alkenyl heteroatom compounds because of its advantages in terms of synthetic efficiency and atom economy.<sup>1</sup> Over the past several decades, considerable efforts have been made in this area and some important alkenyl heteroatom compounds, such as phosphonates,<sup>2</sup> boronates,<sup>3</sup> selenides,<sup>4</sup> sulfides,<sup>5</sup> and nitrogen-<sup>6</sup> and oxygen-containing<sup>7</sup> products, have been obtained *via* the catalytic addition reaction of alkynes. Nevertheless, there is still a great demand for the development of a new and selective catalytic system to offer other important alkenyl heteroatom compounds such as vinyl sulfones. Herein, we report a novel and efficient copper-catalyzed direct hydrosulfonylation of alkynes with arylsulfonic acids for the construction of vinyl sulfones in a one-pot procedure with 100% atom efficiency (eqn (1)).



As extremely valuable alkenyl heteroatom compounds, vinyl sulfones have increasingly attracted synthetic pursuit of chemists, since they can serve as key structural motifs of many

biologically active compounds<sup>8</sup> and versatile building blocks for various organic transformations.<sup>9</sup> Generally, vinyl sulfones are prepared by Knoevenagel condensation of aromatic aldehydes with sulfonylacetic acids,<sup>10</sup> Wittig reaction,<sup>11</sup> Horner-Emmons reaction of carbonyl compounds with sulfonyl phosphoranes,<sup>12</sup> and  $\beta$ -elimination of selenosulfones or halo-sulfones,<sup>13</sup> in which the desired products were usually formed in a mixture of isomers. A number of alternative methods towards vinyl sulfones synthesis have also been developed such as the oxidation of vinyl sulfides with stoichiometric oxidants,<sup>14</sup> the cross-coupling of sulfinate salts with vinyl triflates, vinyl bromides, or alkenyl boronic acids with Cu or Pd catalysts,<sup>15</sup> the addition of  $\text{ArSO}_2\text{X}$  ( $\text{X} = \text{I}, \text{Cl}, \text{SePh}, \text{HgCl}, \text{ONO}_2$ , etc.) to alkenes followed by  $\beta$ -elimination,<sup>16</sup> and the addition of  $\text{ArSO}_2\text{X}$  ( $\text{X} = \text{I}, \text{Cl}$ , etc.), sodium sulfonates or 1,2-bis-(phenylsulfonyl)ethane to alkynes.<sup>17</sup> However, most of these methods suffer from some limitations such as inaccessible starting materials, tedious procedures, relatively harsh reaction conditions, lack of atom economy, or the generation of large amounts of unwanted byproducts. Therefore, the development of mild, convenient, highly-selective and especially atom-economic, environmentally friendly, and resource efficient methods to afford vinyl sulfones is still highly desirable.

The present method of direct addition of arylsulfonic acids to alkynes is realized under mild conditions by employing simple and cheap copper salts as the catalyst, which provides a variety of vinyl sulfones in good to excellent yields and high regio- and stereoselectivity. To the best of our knowledge, this method is the first example of transition-metal-catalyzed direct synthesis of vinyl sulfones from simple and readily available materials with complete atom economy, and does not require the use of any ligand or additive.

Initially, the reaction of phenylacetylene **1a** with benzenesulfonic acid **2a** was performed to examine the catalytic activity of various transition metal complexes including Ru, Pd, Ag, Cu, Fe, Co, Ni, In and Zn salts in DME at 60 °C under  $\text{N}_2$ . As shown in Table 1, among the above metal catalysts screened (entries 1–13), copper salts especially  $\text{Cu(OAc)}_2$  were found to

<sup>a</sup>The Key Laboratory of Life-Organic Analysis, Qufu Normal University, Qufu 273165, China. E-mail: huawang\_qfnu@126.com

<sup>b</sup>Key Laboratory of Pharmaceutical Intermediates and Analysis of Natural Medicine, Qufu Normal University, Qufu 273165, China

<sup>c</sup>Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining 810001, PR China


Table 1 Optimization of the reaction conditions<sup>a</sup>

Entry	Catalyst	Solvent	Yield <sup>b</sup> (%)
1	RuCl <sub>3</sub>	DME	0
2	Pd(OAc) <sub>2</sub>	DME	0
3	In(OAc) <sub>3</sub>	DME	0
4	AgNO <sub>3</sub>	DME	0
5	NiCl <sub>2</sub>	DME	Trace
6	ZnBr <sub>2</sub>	DME	Trace
7	Co(OAc) <sub>2</sub>	DME	Trace
8	FeBr <sub>3</sub>	DME	<10
9	CuBr <sub>2</sub>	DME	82
10	CuCl <sub>2</sub>	DME	90
11	CuI	DME	50
12	Cu(OTf) <sub>2</sub>	DME	75
13	<b>Cu(OAc)<sub>2</sub></b>	<b>DME</b>	<b>94</b>
14	Cu(OAc) <sub>2</sub>	THF	92
15	Cu(OAc) <sub>2</sub>	1,4-Dioxane	65
16	Cu(OAc) <sub>2</sub>	Toluene	51
17	Cu(OAc) <sub>2</sub>	DMSO	26
18	Cu(OAc) <sub>2</sub>	CH <sub>3</sub> OH	44
19	Cu(OAc) <sub>2</sub>	CH <sub>3</sub> CN	0
20	Cu(OAc) <sub>2</sub>	H <sub>2</sub> O	0
21	Cu(OAc) <sub>2</sub>	DME	42 <sup>c</sup>
22	Cu(OAc) <sub>2</sub>	DME	0 <sup>d</sup>
23	—	DME	0

<sup>a</sup> Reaction conditions: **1a** (0.5 mmol), **2a** (0.75 mmol), catalyst (5 mol%), solvent (3 mL), 60 °C, N<sub>2</sub>, 8 h. <sup>b</sup> Isolated yields based on **1a**. <sup>c</sup> Under air. <sup>d</sup> At room temperature.

be the best catalysts for the formation of product **3aa**, and other catalysts such as Pd(OAc)<sub>2</sub>, RuCl<sub>3</sub>, In(OAc)<sub>3</sub>, AgNO<sub>3</sub>, NiCl<sub>2</sub>, ZnBr<sub>2</sub>, Co(OAc)<sub>2</sub>, or FeBr<sub>3</sub> did not or only sluggishly promoted this reaction. The screening of a range of solvents showed that the reaction performed in 1,2-dimethoxyethane (DME) or THF was significantly better than those in 1,4-dioxane, toluene, DMSO, CH<sub>3</sub>OH, CH<sub>3</sub>CN and H<sub>2</sub>O (entries 13–20). The desired product was obtained in only 42% yield under an air atmosphere (entry 21). No conversion was observed when the reaction was performed at room temperature or in the absence of a copper catalyst (entries 22–23).


Under the optimized conditions, the scope of the reaction of various alkynes with arylsulfonic acids was investigated and the results are shown in Table 2. In general, aromatic alkynes which have electron-donating or withdrawing groups on the aryl rings were suitable for this protocol, and the products were obtained in good to excellent yields. Various functionalities such as halogen, cyano, and alkynyl groups were tolerated in this process to afford the corresponding products **3ea–3ja**, which could be employed for further transformations. Moreover, heteroaromatic alkynes such as 3-ethynylpyridine and 3-ethynylthiophene were also compatible with this reaction, providing the corresponding products (**3ka** and **3la**) in 75% and 91% yields, respectively. Notably, internal alkynes such as prop-1-ynylbenzene could also be used in the reaction to give the expected vinyl sulfone **3ma** in moderate yield. Meantime,

Table 2 Scope of the reaction of various alkynes with arylsulfonic acids<sup>a,b</sup>


<sup>a</sup> Reaction conditions: **1** (0.5 mmol), **2** (0.75 mmol), Cu(OAc)<sub>2</sub> (5 mol%), DME (3 mL), 8 h, N<sub>2</sub> (balloon). <sup>b</sup> Isolated yields based on **1**.

1-ethynyl-naphthalene and 1-ethynylcyclohex-1-ene could be used in the reactions to give the expected products (**3na** and **3oa**) in 64% and 91% yields, respectively. Furthermore, the scope of a variety of arylsulfonic acids was examined. In addition to benzenesulfonic acid **2a**, a series of substituted arylsulfonic acids containing either electron-rich or electron-deficient groups were all suitable substrates, and generated the corresponding products in good yields (**3ab–3ae**). In addition, the sterically hindered substituted arylsulfonic acids such as 2-bromobenzenesulfonic acid and 2-(trifluoromethyl)benzenesulfonic acid were also tolerated in this reaction to afford the products in good yields (**3af** and **3ag**).


The feasibility of the gram-scale reaction was investigated by using the model reaction between **1a** and **2a**. It was found that this reaction could afford 2.1 g of **3aa** in 86% yield without any significant loss of its efficiency (eqn (2)). Therefore, this procedure could serve as a practical and efficient protocol to synthesize vinyl sulfones.



It is established that the oxidative addition of sulfinic acids to alkynes leading to  $\beta$ -keto sulfones proceeded *via* a radical process.<sup>18a</sup> When the reaction of **1a** and **2a** was performed in the copper catalytic system under air (dioxygen), in addition to the desired product **3aa**, the  $\beta$ -keto sulfone **10aa** was also obtained in 20% yield (eqn (3)). Therefore, a radical pathway was also supposed to be involved in this hydrosulfonylation reaction.




Furthermore, when TEMPO (2,2,6,6-tetramethyl-1-piperidinyloxy, a well known radical-capturing species) was added into the present reaction system, this hydrosulfonylation process was significantly inhibited (eqn (4)), indicating that this reaction might involve a radical process.



Based on the above information and previous studies,<sup>17-19</sup> we propose a postulated reaction pathway shown in Scheme 1. Firstly, the reaction of sulfinic acid **2** with Cu(OAc)<sub>2</sub> gave the sulfinyl anion **4**, which could be further oxidized by Cu<sup>II</sup> *via* the single electron transfer (SET) process to afford an oxygen-centered radical **5** resonating with sulfonyl radical **6**.<sup>18</sup> Subsequently, the selective addition of sulfonyl radical **6** to alkyne **1** would lead to the formation of reactive vinyl radical **7**, which interacted with Cu<sup>I</sup> species to yield vinyl copper(II) complexes **8**. Finally, the protonation of **8** produced the desired product **3** and regenerated Cu(II) catalyst.<sup>19</sup> The side product **10** might be formed by the isomerization of intermediate **9**, which was generated from vinyl radical **7** in the presence of air (dioxygen) *via* the redox-transfer process.<sup>18a</sup>

In conclusion, a novel and practical protocol of copper-catalyzed direct hydrosulfonylation of alkynes with arylsulfinic



Scheme 1 Postulated reaction pathway.

acids has been developed under mild conditions. It may possess some advantages of cheap catalysts, readily-available starting materials, operation simplicity, high atom economy and reaction selectivity, opening a new door to the construction of vinyl sulfones. Studies of the detailed mechanism of this process and its application are ongoing.

This work was supported by the National Natural Science Foundation of China (no. 21302109, 21302110, and 21375075), the Taishan Scholar Foundation of Shandong Province, the Excellent Middle-Aged and Young Scientist Award Foundation of Shandong Province (BS2013YY019), and the Scientific Research Foundation of Qufu Normal University (BSQD 2012020).

## Notes and references

- For selective reviews, see: (a) F. Alonso, I. P. Beletskaya and M. Yus, *Chem. Rev.*, 2004, **104**, 3079; (b) T. E. Müller, K. C. Hultsch, M. Yus, F. Foubelo and M. Tada, *Chem. Rev.*, 2008, **108**, 3795; (c) B. Godoi, R. F. Schumacher and G. Zeni, *Chem. Rev.*, 2011, **111**, 2937.
- (a) L.-B. Han, C.-Q. Zhao, S. Onozawa, M. Goto and M. Tanaka, *J. Am. Chem. Soc.*, 2002, **124**, 3842; (b) L.-B. Han, C. Zhang, H. Yazawa and S. Shimada, *J. Am. Chem. Soc.*, 2004, **126**, 5080.
- (a) C. Gunanathan, M. Hölscher, F. Pan and W. Leitner, *J. Am. Chem. Soc.*, 2012, **134**, 14349; (b) H. Yoshida, S. Kawashima, Y. Takemoto, K. Okada, J. Ohshita and K. Takaki, *Angew. Chem., Int. Ed.*, 2012, **51**, 235.
- (a) I. Kamiya, E. Nishinaka and A. Ogawa, *J. Org. Chem.*, 2005, **70**, 696; (b) S. Kawaguchi, M. Kotani, S. Atobe, A. Nomoto, M. Sonoda and A. Ogawa, *Organometallics*, 2011, **30**, 6766.
- (a) A. D. Giuseppe, R. Castarlenas, J. J. Pérez-Torrente, M. Crucianelli, V. Polo, R. Sancho, F. J. Lahoz and L. A. Oro, *J. Am. Chem. Soc.*, 2012, **134**, 8171; (b) S. N. Riduan, J. Y. Ying and Y. Zhang, *Org. Lett.*, 2012, **14**, 1780.
- (a) Y. Fukumoto, H. Asai, M. Shimizu and N. Chatani, *J. Am. Chem. Soc.*, 2007, **129**, 13792; (b) S. Yudha, Y. Kuninobu and K. Takai, *Org. Lett.*, 2007, **9**, 5609.
- (a) B. C. Chary and S. Kim, *J. Org. Chem.*, 2010, **75**, 7928; (b) M. Kawatsura, J. Namioka, K. Kajita, M. Yamamoto, H. Tsuji and T. Itoh, *Org. Lett.*, 2011, **13**, 3285.
- (a) M. Uttamchandani, K. Liu, R. C. Panicker and S. Q. Yao, *Chem. Commun.*, 2007, 1518; (b) J. J. Reddick, J. Cheng and W. R. Roush, *Org. Lett.*, 2003, **5**, 1967; (c) G. Wang, U. Mahesh, G. Y. J. Chen and S. Q. Yao, *Org. Lett.*, 2003, **5**, 737; (d) D. C. Meadows, T. Sanchez, N. Neamati, T. W. North and J. Gervay-Hague, *Bioorg. Med. Chem.*, 2007, **15**, 1127; (e) B. A. Frankel, M. Bentley, R. G. Kruger and D. G. McCafferty, *J. Am. Chem. Soc.*, 2004, **126**, 3404.
- (a) H. Kumamoto, K. Deguchi, T. Wagata, Y. Furuya, Y. Odanaka, Y. Kitade and H. Tanaka, *Tetrahedron*, 2009,

- 65, 8007; (b) Q. Zhu and Y. Lu, *Org. Lett.*, 2009, **11**, 1721; (c) G. Pandey, K. N. Tiwari and V. G. Puranik, *Org. Lett.*, 2008, **10**, 3611; (d) M. N. Noshi, A. El-awa, E. Torres and P. L. Fuchs, *J. Am. Chem. Soc.*, 2007, **129**, 11242; (e) D. L. J. Clive, Z. Li and M. Yu, *J. Org. Chem.*, 2007, **72**, 5608; (f) D. J. Wardrop and J. Fritz, *Org. Lett.*, 2006, **8**, 3659; (g) S. Sulzer-Mossé, A. Alexakis, J. Mareda, G. Bollot, G. Bernardinelli and Y. Filinchuk, *Chem.–Eur. J.*, 2009, **15**, 3204; (h) A. López-Pérez, R. Robles-Machín, J. Adrio and J. C. Carretero, *Angew. Chem., Int. Ed.*, 2007, **46**, 9261.
- 10 (a) S. Chodroff and W. F. Whitmore, *J. Am. Chem. Soc.*, 1950, **72**, 1073; (b) D. A. R. Happer and B. E. Steenson, *Synthesis*, 1980, 806.
- 11 (a) M. Mikolajczyk, W. Perlikowska, J. Omelanczuk, H. Cristau and A. Perraud-Darcy, *J. Org. Chem.*, 1998, **63**, 9716; (b) J. H. van Steenis, J. J. G. S. van Es and A. van der Gen, *Eur. J. Org. Chem.*, 2000, 2787.
- 12 I. C. Popoff and J. L. Denver, *J. Org. Chem.*, 1969, **34**, 1128.
- 13 (a) R. A. Gancarz and J. L. Kice, *Tetrahedron Lett.*, 1980, **21**, 4155; (b) M. Asscher and D. Vofsi, *J. Chem. Soc., Perkin Trans. 1*, 1972, 1543; (c) P. B. Hopkin and P. L. Fuchs, *J. Org. Chem.*, 1978, **43**, 1208.
- 14 (a) M. Kirihara, J. Yamamoto, T. Noguchi and Y. Hirai, *Tetrahedron Lett.*, 2009, **50**, 1180; (b) Q. Xue, Z. Mao, Y. Shi, H. Mao, Y. Cheng and C. Zhu, *Tetrahedron Lett.*, 2012, **53**, 1851.
- 15 (a) G. W. Kabalka and S. K. Guchhait, *Tetrahedron Lett.*, 2004, **45**, 4021; (b) S. Cacchi, G. Fabrizi, A. Goggiamani, L. M. Parisi and R. Bernini, *J. Org. Chem.*, 2004, **69**, 5608; (c) F. Huang and R. A. Batey, *Tetrahedron*, 2007, **63**, 7667; (d) D. C. Reeves, S. Rodriguez, H. Lee, N. Hahhad, D. Krishnamurthy and C. H. Senanayake, *Tetrahedron Lett.*, 2009, **50**, 2870.
- 16 (a) L. K. Liu and K.-Y. Jen, *J. Org. Chem.*, 1980, **45**, 406; (b) C. Nájera, B. Baldó and M. Yus, *J. Chem. Soc., Perkin Trans. 1*, 1988, 1029; (c) V. Nair, A. Augustine, T. G. George and L. G. Nair, *Tetrahedron Lett.*, 2001, **42**, 6763; (d) V. Nair, A. Augustine and T. D. Suja, *Synthesis*, 2002, 2259; (e) T. G. Back and S. Collins, *J. Org. Chem.*, 1981, **46**, 3249; (f) V. Nair, A. Augustine, S. B. Panicker, T. D. Suja and S. Mathai, *Res. Chem. Intermed.*, 2003, **29**, 213; (g) H. Qian and X. Huang, *Synlett*, 2001, 1913.
- 17 (a) X. Liu, X. Duan, Z. Pan, Y. Han and Y. Liang, *Synlett*, 2005, 1752; (b) N. Taniguchi, *Synlett*, 2011, 1308; (c) N. Taniguchi, *Synlett*, 2012, 1245; (d) R.-J. Song, Y. Liu, Y.-Y. Liu and J.-H. Li, *J. Org. Chem.*, 2011, **76**, 1001.
- 18 (a) Q. Lu, J. Zhang, G. Zhao, Y. Qi, H. Wang and A. W. Lei, *J. Am. Chem. Soc.*, 2013, **135**, 11481; (b) Q. Lu, J. Zhang, F. Wei, Y. Qi, H. Wang, Z. Liu and A. W. Lei, *Angew. Chem., Int. Ed.*, 2013, **52**, 7156.
- 19 (a) A. Kondoh, H. Yorimitsu and K. Oshima, *J. Am. Chem. Soc.*, 2007, **129**, 4099; (b) S. Kim and P. H. Lee, *J. Org. Chem.*, 2012, **77**, 215.